JA J

JPRS: 3120

24 March 1960

MAIN FILE

THE PROBLEM OF HYDROGEN ISOTOPE EXCHANGE IN

SATURATED HYDROCARBONS

(USSR)

[Translation]

्रकार व्यक्ति । व्यक्ति व्यक्ति । अञ्चल व्यक्ति व्यक्ति व्यक्ति ।

Approved for Public Release
Distribution Unlimited

Continue Co

Distributed by:

OFFICE OF TECHNICAL SERVICES U.S. DEPARTMENT OF COMMERCE WASHINGTON 25, D. C. (Price: \$0.50)

20000104 068

U. S. JOINT FUBLICATIONS RESEARCH SERVICE
205 EAST 42nd STREET, SUITE 300
NEW YORK 17, N. Y.

Reproduced From Best Available Copy

DEED OFMERTY LINES OF THE 2

FOREWORD

This publication was prepared under contract by the UNITED STATES JOINT PUBLICATIONS RE-SEARCH SERVICE, a federal government organization established to service the translation and research needs of the various government departments.

A 1973 COTAGE MATERIA TOO.

SO SETON OF ON NOT DO FOREIA

Foreign to noblect to be

JFRS: 3120 CSO: 3481-I

THE PROBLEM OF HYDROGEN ISOTOPE EXCHANGE IN

SATURATED HYDROCARBONS

(USSR)

Ya. M. Varshavskiy S. E. Vaysberg M. G. Lozhkina

In conducting hydrogen isotope exchange reactions in heavy water, as is well known, the hydrogen atoms in C-H hydrocarbon bonds are either exchanged with great difficulty or practically are not quite changed to deuterium. A. I. Brodskiy /1/ figured that it was bonded in the absence of free electron pairs from carbon. A. I. Shatenshteyn and co-workers /2-4/ showed, that a hydrogen exchange in C-H bonds is sharply accelerated if the reaction is conducted in such mediums as: deuteroammonia, deuterium bromide or deuterium flouride. However even in such strongly heavy dydrogen mediums, as ND₃ and DBr, hydrogen atoms of saturated hyhyocarbons do not exchange.

The problem of hydrogen isotope exchange in saturated hydrocarbons has its own history. Taylor, Morikava and Benedict [5], who succeeded in accomplishing the hydrogen exchange in a gas phase between methane and deuterium by catalyzation with agitated copper atoms, first observed such an exchange in 1935. Later [6] an exchange between methane and ethane on the one hand and atomized deuterium on the other was accomplished. The catalytic action of several metals and oxides on a hydrogen exchange in C-H bonds was studied in a series of works. By this it was shown that the exchange occurs comparatively easily in olefin and aromatic hydrocarbons, whereas in the case of saturated hydrocarbons even in very active catalyzers at high temperature it does not take place or it takes place with great difficulty.

Ingold, Raisin, and Wilson observed the hydrogen exchange of saturated hydrocarbons in heavy hydrogenized sulfuric acid for the first time in 1936 \int 7. Later, the exchange of hydrocarbons in deuterosulfuric acid was studied in more detail \int 8-10/. It was pointed out that in the case of hydrocarbons, not containing a tertiary carbon atom, the exchange is not noted, whereas a complete exchange is observed in the presence of even one tertiary C-atom in the molecule. This fact was

explained by the exidation of the tertiary carbon atom_by sulfuric acid with the following exchange in the carbonium ion /9,10/ or olefin /8/), being formed as a result of the exidation. Since in the exchange process the complete exidation of hydrocarbon was not observed, the hypothesis was expressed, according to which the carbonium center (or double bond) is transferred both inside the molecule and from one mulecule to the other according to the chain mechanism. Thus the presence of a tertiary carbon atom, which may easily be exidized is acknowledge the necessary condition of an exchange reaction in deuterosulfuric acid and also the necessity of the very act of exidation, being the beginning of the chain.

Data [2,11] on the insignificant introduction of deuterium into n-heptane, cyclohexane and decalin molecules by condunting the exchange in a potassium amide solution in liquid heavy hydrogenized ammonia is also available in the literature.

the medium of liquid deuterium bromide in the presence of AIBr3. It was shown, that the exchange reaction in such a medium occurs comparatively easy and in the course of 60 hours at room temperature equilibrium may be attained practically.* (* A short reference of the results of these tests were made in the footnote to the article / 3 / and also in the summary / 2 /.) For greater certainty, besides the tests on the direct exchange, tests on the reverse exchange also were carried out, for which at first cyclohexane was heavy hydrogenized and then dissolved in an AIBr3 solution in hydrogen bromide, not containing deuterium, and the introduction of deuterium from cyclohexane into hydrogen bromide was tested.** (** The absence of the isomerization of cyclohexane in methylcyclopentane was checked out in our tests by the invariability of the boiling temperature of cyclohexane 81.0°c) up to and after the exchange.)

That case, wherein the hydrogen exchange in saturated hydrocarbons, containing a tertairy carbon atom, must not certainly pass through the oxidation stage, is confirmed by the tests on the isotope exchange in the medium of deuterium flouride.

Earlier [4] it was shown that such saturated hydrocarbons as methylcyclohexane and decalin, exchange their own hydrogen atoms sufficiently easy in a medium of liquid deuterium flouride, and in the presence of BF3 the exchange may also be successfully observed even in cyclohexane. Tests, which we conducted recently with the application of specially prepared pure deuterium flouride without oxidizers, confirmed the earlier obtained results [4]. At room temperature in the course of 190 hours 11 hydrogen atoms were exchanged in methylcyclohexane and in the course of 760 hours 13 atoms exchanged.

The obtained results witnessed the fact that AlBr3 solutions in hydrogen bromide and also hydrogen flouride and more so BF3 solutions in HF* may be used with success in the capacity of a medium for conducting systematic tests of hydrogen isotope exchange in saturated hydrocarbons (* The accleration of the hydrogen exchange in hydrocarbons in a medium of deuterium flouride ought to be expected also by the dissolving of such flourides, as AsF5 and SbF5 $\int 12/$ in it.)

These results also witness, first, the fact, that an exchange in saturated hydrocarbons must not certainly be accompanied by their oxidation, since hydrogen bromide, hydrogen flouride and also AlBr3 and BF3 do not possess oxidizing characteristics, and second, the fact, that the presence of a tertiary carbon atom in the molecule is not compulsory for the exchange.

The problem of the mechanism of our observed fast hydrogen exchange in saturated hydrocarbons arouses the principal interest. This is connected with the fact that in saturated hydrocarbons, as is well known, easily polarized 77 -bonds are lacking, at the expense of which intermediate complexes with proton donors or carbonium ions may be produced comparatively easy. Not excluded was the fact, that our studied exchange reactions pass through the stage of the joining of deutron to the saturated hydrocarbon molecule from the positively formed charged ion; it is also possible that the process of deuterium introduction in the form of deutron, and hydrogen loss in the form of a proton occurs in a single act according to the Walden inversion type mechanism. In conclusion we express thanks to the chief of the laboratory, professor A. I. Shatenshteyn for valuable advice.

L. A. Karpov Physical-Chemistry Institute Published 15 Jan. 1955.

Literature

- 1. A. I. Brodskiy, <u>Izv. AN SSSR</u>, <u>OXH Otdeleniya Khimicheskika Nauk</u>
 <u>Izvestiya Akademii Nauk, SSSR</u>, <u>/Proceedings</u> of the Academy of Science,
 USSR, Department of Chemical Science, 1, 3, 1949.
- 2. A. I. Shatenshteyn, <u>Uspekhi Khimii</u> ∫Successes of Chemistry, 21, 914, 1952.
- 3. A. I. Shatenshteyn, Ya. M. Varshavskiy, <u>DAN</u> Doklady Akademii Nauk-Reports of the Academy of Science/85, 157, 1952.
- 4. Ya. M. Varshavskiy and A. I. Shatenshteyn, DAN, 95, 297, 1954.
- 5. H. S. Taylor, K. Morikava, W. S. Benedict, <u>Journ. Amer. Chem. Soc.</u>, 57, 383, 1935.

- E. W. R. Steacie, N. W. E. Phyllips, Journ. Chem. Phys. 4, 461, 1936.
- C. K. Ingold, C. G. Raisin, C. L. Wilson, <u>Journ. Chem. Soc.</u> (L), 1643, 1936.
 R. L. Burwell, G. S. Gordon, <u>Journ, Amer. Chem. Soc.</u>, 70, 3128, 1948
- 71, 2355, 1949.
- V. N. Setkina, D. N. Kursanov, O. D. Sterligov, A. L. Liberman, DAN, 85, 1045, 1952. and the artificial property and an array of the second transfer of the
- 10. J. W. Otvos, D. P. Stenvenson, C. D. Wagner, O. Beek, Journ., Amer. Chem. Soc., 74, 3269, 1952.
- 11. A. I. Shatenshteyn, L. N. Vasil'eva, N. M. Dykhno, I. A. Izrailevich, DAN, 85, 381, 1952, a secretary of the real transfer and the metaling believed
- 12. L. F. Audrieth, J. Kleinbert, "Non-aqueous solvents," 1953 (Chapter 10) (Russian translation Ordit i Kleynberg, Non-aqueous solvents, IIL /Izdatel'stvo Inostrannoy Literatury/, 1955, Chapter 10)/. Foreign Literature Publishing House en de la companya de la co

Burger : Francisco Carlos Carl

Company by Local St.

ung und Punkan bei gestern Dichte wilder zu der Anderschieß in der

The control of the second of t

gand and the second of the sec

The state of the s

and the second of the second o

ति । हैं। ते स्थापित करा करा करा है। ते हैं ते हुए हैं हुई स्थापित स्थापित स्थाप है। स्थापित है स्थापित है

5810, which is the street of t

STARRED BURNES

_ 4 _